

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Rsrll
Catalog Number: R0501S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10173034
Expiration Date: 12/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.15% Triton® X-100, 200 μg/ml BSA (pH 7.4 @ 25°C)

Specification Version: PS-R0501S/L v2.0

RsrII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0501SVIAL	Rsrll	10173033	Pass	
B6004SVIAL	rCutSmart™ Buffer	10173161	Pass	

Assay Name/Specification	Lot # 10173034
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of Lambda DNA and a minimum of 50 units of RsrII incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with RsrII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with RsrII.	
Protein Purity Assay (SDS-PAGE)	Pass
RsrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
detection.	
Endonuclease Activity (Nicking)	Pass
A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and	
a minimum of 50 units of RsrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
conversion to the micked form as determined by againse gerelectrophoresis.	



R0501S / Lot: 10173034

Page 1 of 2

Assay Name/Specification	Lot # 10173034
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of RsrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń

Production Scientist

13 Dec 2022

Michael Tonello

Packaging Quality Control Inspector

14 Dec 2022



R0501S / Lot: 10173034

Page 2 of 2