

## New England Biolabs Certificate of Analysis

**Product Name:** *EaeI*  
**Catalog Number:** *R0508L*  
**Concentration:** *5,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Packaging Lot Number:** *10080204*  
**Expiration Date:** *07/2022*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*  
**Specification Version:** *PS-R0508S/L v1.0*

EaeI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0508LVIAL	EaeI	10080203	Pass
B7204SVIAL	CutSmart® Buffer	10078756	Pass

Assay Name/Specification	Lot # 10080204
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 25 units of EaeI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with EaeI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EaeI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 15 Units of EaeI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
17 Sep 2020



Michael Tonello  
Packaging Quality Control Inspector  
17 Sep 2020