

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	PfIMI
Catalog Number:	R0509S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10098186
Expiration Date:	01/2023
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0509S/L v1.0

PfIMI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0509SVIAL	PfIMI	10098187	Pass	
B7203SVIAL	NEBuffer™ 3.1	10092686	Pass	

Assay Name/Specification	Lot # 10098186
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled pNEB193 DNA and a minimum of 50 units of PfIMI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) PfIMI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 μl reaction in NEBuffer 3.1 containing 1 μg of Lambda DNA and a minimum of 30 Units of PfIMI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with PfIMI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with PfIMI.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10098186
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and	
double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of PfIMI incubated for	
4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 10 Feb 2021

Josh Hersey

Packaging Quality Control Inspector 10 Feb 2021

