240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: AflII

Catalog #: R0520S/L
Concentration: 20,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 RF I DNA in 1 hour at 37°C in a total

reaction volume of 50 µl.

Lot #:0431502Assay Date:02/2015Expiration Date:02/2017Storage Temp: $-20 \, ^{\circ}$ C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R0520S/L v1.0
Effective Date: 26 Nov 2014

| Assay Name/Specification (minimum release criteria)  | Lot #0431502 |
|--|--------------|
| <b>Blue-White Screening (Terminal Integrity)</b> - A sample of Litmus28 vector linearized with a 10-fold excess of AfIII, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.  | Pass         |
| Endonuclease Activity (Nicking) - A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of supercoiled pBR322 DNA and a minimum of 100 Units of AfIII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.                          | Pass         |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of AflII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 5-fold over-digestion of PhiX174 DNA with AfIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AfIII.   | Pass         |
| Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of PhiX174 DNA and a minimum of 200 Units of AfIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.           | Pass         |







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (#R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

Authorized by Derek Robinson 26 Nov 2014







Inspected by John Greci 23 Feb 2015