

DISCOVERY 240 County Road
GENUINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: AfIII

Catalog Number: R0520S

Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of PhiX174 RF I DNA in 1 hour at 37°C in a total reaction volume of

50 μl.

Packaging Lot Number: 10073397 Expiration Date: 12/2021 Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0520S/L v1.0

AfIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0520SVIAL	AfIII	10061314	Pass	
B7204SVIAL	CutSmart® Buffer	10074633	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10071081	Pass	

Assay Name/Specification	Lot # 10073397
Blue-White Screening (Terminal Integrity) A sample of Litmus28 vector linearized with a 10-fold excess of AfIII, religated	Pass
and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 100 Units of AfIII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart [™] Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of AfIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of PhiX174 DNA with AfIII, >95% of the DNA fragments	Pass



R0520S / Lot: 10073397

Page 1 of 2

Assay Name/Specification	Lot # 10073397
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AfIII.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of PhiX174 DNA and a minimum of	
200 Units of AfIII incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang Production Scientist

18 Jun 2020

Michael Tonello

Packaging Quality Control Inspector

18 Jun 2020



R0520S / Lot: 10073397

Page 2 of 2