

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Xcml
Catalog Number: R0533S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 1009735 Expiration Date: 01/2023 Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0533S/L v1.0

| Xcml Component List    |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0533SVIAL             | Xcml                  | 10097355   | Pass                 |  |
| B7202SVIAL             | NEBuffer™ 2.1         | 10090560   | Pass                 |  |

| Assay Name/Specification  | Lot # 10097356 |
|---|----------------|
| Non-Specific DNase Activity (16 hour) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 5 Units of Xcml incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme. | Pass           |
| Ligation and Recutting (Terminal Integrity)  After a 10-fold over-digestion of Lambda DNA with XcmI, ~25% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with XcmI.  | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 2.1 containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of Xcml incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.   | Pass           |

This product has been tested and shown to be in compliance with all specifications.



R0533S / Lot: 10097356 Page 1 of 2 One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Production Scientist

16 Feb 2021

Michael Tonello

Packaging Quality Control Inspector

16 Feb 2021



R0533S / Lot: 10097356

Page 2 of 2