

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Bsal
Catalog Number:	R0535S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10091140
Expiration Date:	08/2022
Storage Temperature:	-20°C
Storage Conditions:	200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0535S/L v2.0

Bsal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0535SVIAL	Bsal	10081424	Pass	
B7204SVIAL	CutSmart® Buffer	10089397	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084973	Pass	

Assay Name/Specification	Lot # 10091140
Protein Purity Assay (SDS-PAGE) Bsal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of pXba DNA and a minimum of 50 units of Bsal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pXba DNA with Bsal, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Bsal.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Bsal incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10091140
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 10 units of Bsal incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 18 Nov 2020

Michae

Michael Tonello Packaging Quality Control Inspector 18 Nov 2020

