

New England Biolabs Certificate of Analysis

Product Name: BbsI
Catalog Number: R0539L
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r2.1 in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10197907
Expiration Date: 06/2025
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml rAlbumin, (pH 7.4 @ 25°C)
Specification Version: PS-R0539S/L v3.0

BbsI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0539LVIAL	BbsI	10197906	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10184699	Pass
B6002SVIAL	NEBuffer™ r2.1	10156432	Pass


Assay Name/Specification	Lot # 10197907
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of BbsI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of BbsI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of BbsI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of a mixture of single and	Pass

Assay Name/Specification	Lot # 10197907
<p>double-stranded [³H] E. coli DNA and a minimum of 50 units of BbsI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	
<p>Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 µl of BbsI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 µl of BbsI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BbsI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours hours at 25°C. Of these ligated fragments, >95% can be recut with BbsI.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BbsI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours hours at 25°C. Of these ligated fragments, >95% can be recut with BbsI.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and a minimum of 50 units of BbsI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and a minimum of 50 units of BbsI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of BbsI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of BbsI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results</p>	Pass

Assay Name/Specification	Lot # 10197907
are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
11 Jul 2023



Michael Tonello
Packaging Quality Control Inspector
17 Jul 2023