

## New England Biolabs Certificate of Analysis

**Product Name:** BbsI  
**Catalog Number:** R0539S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r2.1 in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10197908  
**Expiration Date:** 06/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml rAlbumin, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0539S/L v3.0

BbsI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0539SVIAL	BbsI	10197903	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10193044	Pass
B6002SVIAL	NEBuffer™ r2.1	10173667	Pass


Assay Name/Specification	Lot # 10197908
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of BbsI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of BbsI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of BbsI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of a mixture of single and	Pass

Assay Name/Specification	Lot # 10197908
<p>double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 50 units of BbsI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	
<p><b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 µl of BbsI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 µl of BbsI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with BbsI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours hours at 25°C. Of these ligated fragments, &gt;95% can be recut with BbsI.</p>	<b>Pass</b>
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with BbsI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours hours at 25°C. Of these ligated fragments, &gt;95% can be recut with BbsI.</p>	<b>Pass</b>
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and a minimum of 50 units of BbsI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and a minimum of 50 units of BbsI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of BbsI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of BbsI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10197908
are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is $\leq 1$ E. coli genome.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
11 Jul 2023



Josh Hersey  
Packaging Quality Control Inspector  
03 Aug 2023