

New England Biolabs Certificate of Analysis

Product Name: *BspEI*
Catalog Number: *R0540L*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (dam -) in NEBuffer r3.1 in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10253160*
Expiration Date: *07/2026*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)*
Specification Version: *PS-R0540S/L v3.0*


BspEI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0540LVIAL	BspEI	10248163	Pass
B6003SVIAL	NEBuffer™ r3.1	10237086	Pass

Assay Name/Specification	Lot # 10253160
Blue-White Screening (Terminal Integrity) A sample of LITMUS38i vector linearized with a 10-fold excess of BspEI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of BspEI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of BspEI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda dam- DNA and 1 µl of BspEI incubated for 15 minutes at 37°C results in complete digestion as determined	Pass

Assay Name/Specification	Lot # 10253160
by agarose gel electrophoresis.	
<p>Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda dam- DNA with BspEI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BspEI.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda dam- DNA and a minimum of 50 units of BspEI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) BspEI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of BspEI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
29 Jul 2024



Josh Hersey
Packaging Quality Control Inspector
08 Aug 2024