

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	BspEl
Catalog Number:	R0540S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (dam) in 1 hour at 37°C in a total reaction volume of 50 μl.
Lot Number:	10012829
Expiration Date:	06/2020
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version:	PS-R0540S/L v1.0

BspEl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0540SVIAL	BspEl	10012830	Pass	
B7203SVIAL	NEBuffer™ 3.1	0541804	Pass	

Assay Name/Specification	Lot # 10012829
Blue-White Screening (Terminal Integrity) A sample of LITMUS38i vector linearized with a 10-fold excess of BspEI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 μ I reaction in NEBuffer 3.1 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 10 Units of BspEI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of BspEI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda dam- DNA with BspEI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10012829
fragments, >95% can be recut with BspEI.	
Non-Specific DNase Activity (16 Hour) A 50 μI reaction in NEBuffer 3.1 containing 1 μg of Lambda dam- DNA and a minimum of	Pass
50 units of BspEI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist 18 Jun 2018

Josh Hersey

Packaging Quality Control Inspector 19 Jun 2018

