

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	AfIII
Catalog Number:	R0541S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer™ r3.1 in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number:	10158866
Expiration Date:	08/2024
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 500 mM NaCl ,1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 200 μg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version:	PS-R0541S/L v2.0

AfIIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0541SVIAL	AfIII	10158864	Pass	
B6003SVIAL	NEBuffer™ r3.1	10146825	Pass	

Assay Name/Specification	Lot # 10158866
Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of Lambda DNA with AfIIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AfIIII.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and 1 µl of AfIIII incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and a minimum of 30 units of AfIIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) AfIIII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10158866 Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of AfIIII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of AfIIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 08 Sep 2022

Erin Varney

Packaging Quality Control Inspector 08 Sep 2022

