

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: BcoDl
Catalog Number: R0542S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10198131
Expiration Date: 06/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

200 $\mu g/ml$ rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0542S v2.0

BcoDI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0542SVIAL	BcoDI	10196886	Pass	
B6004SVIAL	rCutSmart™ Buffer	10193042	Pass	

Assay Name/Specification	Lot # 10198131
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and	Pass
double-stranded [³H] E. coli DNA and a minimum of 100 units of BcoDI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of BcoDI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BcoDI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BcoDI.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 30 units of BcoDl incubated for 16 hours at 37°C results in a DNA pattern free of	Pass



R0542S / Lot: 10198131

Page 1 of 2

Assay Name/Specification	Lot # 10198131	
detectable nuclease degradation as determined by agarose gel electrophoresis.		
Protein Purity Assay (SDS-PAGE) BcoDI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass	
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of BcoDI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun \
Production Scientist

27 Jun 2023



Packaging Quality Control Inspector

14 Jul 2023



R0542S / Lot: 10198131

Page 2 of 2