

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	DpnII
Catalog Number:	R0543L
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of 50 μl.
Lot Number:	10014860
Expiration Date:	07/2020
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version:	PS-R0543S/L v1.0

DpnII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0543LVIAL	DpnII	10014861	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	0261805	Pass	
B0543SVIAL	NEBuffer™ DpnII	0091805	Pass	

Assay Name/Specification	Lot # 10014860
<b>Protein Purity Assay (SDS-PAGE)</b> DpnII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of Lambda dam- DNA and a minimum of 100 units of DpnII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of DpnII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda dam- DNA with DpnII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10014860
fragments, >95% can be recut with DpnII.	
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of DpnII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

Stephonie Onet

Stephanie Cornelio Production Scientist 09 Jul 2018

Josh Hersey

Packaging Quality Control Inspector 11 Jul 2018

