

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Dpnll
Catalog Number:	R0543L
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10107052
Expiration Date:	05/2023
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version:	PS-R0543S/L v1.0

DpnII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0543LVIAL	DpnII	10107053	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10093123	Pass	
B0543SVIAL	NEBuffer™ DpnII	10095951	Pass	

Assay Name/Specification	Lot # 10107052
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of DpnII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of DpnII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda dam- DNA with DpnII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DpnII.	Pass
Protein Purity Assay (SDS-PAGE) DpnII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10107052
detection.	
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer DpnII containing 1 µg of Lambda dam- DNA and a minimum of 100 units of DpnII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Pepholog

Penghua Zhang Production Scientist 04 Jun 2021

Michael Tonello Packaging Quality Control Inspector 04 Jun 2021

