240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: PacI

Catalog #: R0547S/L
Concentration: 10,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193 DNA in 1 hour at 37°C in a total reaction

volume of 50  $\mu$ l.

 Lot #:
 0661701

 Assay Date:
 01/2017

 Expiration Date:
 1/2019

 Storage Temp:
 -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R0547S/L v1.0
Effective Date: 19 Feb 2015

Assay Name/Specification (minimum release criteria)	Lot #0661701
<b>Blue-White Screening (Terminal Integrity)</b> - A sample of pNEB193 vector linearized with a 10-fold excess of PacI, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Endonuclease Activity (Nicking)</b> - A 50 μl reaction in CutSmart <sup>™</sup> Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 30 Units of PacI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of PacI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of pNEB193 DNA with PacI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PacI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of pNEB193 DNA and a minimum of 100 units of PacI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> - PacI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

Authorized by Derek Robinson 19 Feb 2015







Inspected by Anthony Francis 11 Jan 2017