

New England Biolabs Certificate of Analysis


Product Name: *Pacl*
Catalog Number: *R0547S*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*
Lot Number: *10045000*
Expiration Date: *04/2021*
Storage Temperature: *-20°C*
Storage Conditions: *200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version: *PS-R0547S/L v1.0*

Pacl Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0547SVIAL	Pacl	10038547	Pass
B7204SVIAL	CutSmart® Buffer	10043347	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10038713	Pass

Assay Name/Specification	Lot # 10045000
Blue-White Screening (Terminal Integrity) A sample of pNEB193 vector linearized with a 10-fold excess of Pacl, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of Pacl incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Pacl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pNEB193 DNA with Pacl, ~75% of the DNA fragments	Pass

Assay Name/Specification	Lot # 10045000
<p>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PaeI.</p>	
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pNEB193 DNA and a minimum of 100 units of PaeI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) PaeI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Anthony Francis
Production Scientist
06 Mar 2019



Mary Conlon
Packaging Quality Control Inspector
19 Jun 2019