

## New England Biolabs Certificate of Analysis

**Product Name:** Pacl  
**Catalog Number:** R0547S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10119695  
**Expiration Date:** 02/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0547S/L v1.0

Pacl Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0547SVIAL	Pacl	10097999	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10114156	Pass
B6004SVIAL	rCutSmart™ Buffer	10118973	Pass

Assay Name/Specification	Lot # 10119695
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of pNEB193 DNA with Pacl, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Pacl.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pNEB193 DNA and a minimum of 100 units of Pacl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Pacl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	Pass

Assay Name/Specification	Lot # 10119695
a minimum of 30 Units of PacI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
<p><b>Blue-White Screening (Terminal Integrity)</b> A sample of pNEB193 vector linearized with a 10-fold excess of PacI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in &lt;1% white colonies.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> PacI is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
21 Sep 2021



Michael Tonello  
Packaging Quality Control Inspector  
21 Sep 2021