

DISCOVERY 240 County Road
GENUINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sapl
Catalog Number: R0569L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10007472
Expiration Date: 06/2020
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R0569S/L v1.0

Sapl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0569LVIAL	Sapl	10007473	Pass	
B7204SVIAL	CutSmart® Buffer	10010632	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	0261805	Pass	

Assay Name/Specification	Lot # 10007472
Protein Purity Assay (SDS-PAGE)	Pass
Sapl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 30 units of Sapl incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled LITMUS38i DNA	
and a minimum of 10 units of Sapl incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
conversion to the nicked form as determined by agarose gerelectrophoresis.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	
30 Units of Sapl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
detectable fluciease degradation as determined by againse generatiophoresis.	



R0569L / Lot: 10007472

Page 1 of 2

Assay Name/Specification	Lot # 10007472
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with Sapl, >95% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with SanI	

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso Production Scientist

12 Jun 2018

Michael Tonello

Packaging Quality Control Inspector

27 Jun 2018

