

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: BsmFl
Catalog Number: R0572L
Concentration: 2,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA in 1 hour at 65°C in a total reaction volume of 50 μl.

Lot Number: 10053625
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0572S/L v1.0

| BsmFI Component List   |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0572LVIAL             | BsmFI                 | 10053626   | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer      | 10046087   | Pass                 |  |

| Assay Name/Specification   | Lot # 10053625 |
|--|----------------|
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 2 Units of BsmFl incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 Units of BsmFl incubated for 4 hours at 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.               | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 20 units of BsmFl incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.               | Pass           |
| Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of pBR322 DNA with BsmFI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsmFI.   | Pass           |



R0572L / Lot: 10053625

Page 1 of 2

This product has been tested and shown to be in compliance with all specifications.

Production Scientist

09 Aug 2019

ayiviinichieilo

Packaging Quality Control Inspector

27 Aug 2019

R0572L / Lot: 10053625

Page 2 of 2