

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Fsel
Catalog Number: R0588S
Concentration: 2,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10173228
Expiration Date: 08/2023
Storage Temperature: -80°C

Storage Conditions: 10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.5 % Tween®

20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R0588S/L v3.0

Fsel Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0588SVIAL	Fsel	10162512	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10162784	Pass	
B6004SVIAL	rCutSmart™ Buffer	10165689	Pass	

Assay Name/Specification	Lot # 10173228
Protein Purity Assay (SDS-PAGE) Fsel is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 Units of Fsel incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 10 units of Fsel incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBC4 DNA and a minimum of 10 units of Fsel incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



R0588S / Lot: 10173228

Page 1 of 2

Assay Name/Specification	Lot # 10173228
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pBC4 DNA with Fsel, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95%	
can be recut with Fsel.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun \
Production Scientist

30 Aug 2022

Michael Tonello

Packaging Quality Control Inspector

29 Nov 2022



R0588S / Lot: 10173228

Page 2 of 2