

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Mfel
Catalog Number: R0589S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10088518
Expiration Date: 11/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0589S/L v2.0

Mfel Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0589SVIAL	Mfel	10088519	Pass	
B7204SVIAL	CutSmart® Buffer	10089402	Pass	

Assay Name/Specification	Lot # 10088518
Blue-White Screening (Terminal Integrity) A sample of LITMUS38i vector linearized with a 10-fold excess of Mfel, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene	Pass
results in <1% white colonies.	
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of Mfel incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 10 units of Mfel incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with MfeI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with MfeI.	Pass



R0589S / Lot: 10088518

Page 1 of 2

Assay Name/Specification	Lot # 10088518
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μl reaction in CutSmart™ Buffer containing 1 μg of Lambda DNA and a minimum of	
30 Units of Mfel incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

28 Nov 2020

Josh Hersey

Packaging Quality Control Inspector

28 Nov 2020

