

New England Biolabs Certificate of Analysis

Product Name: HpyCH4III
Catalog Number: R0618L
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10251735
Expiration Date: 03/2026
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version: PS-R0618S/L v3.0


HpyCH4III Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0618LVIAL	HpyCH4III	10233799	Pass
B6004SVIAL	rCutSmart™ Buffer	10249064	Pass

Assay Name/Specification	Lot # 10251735
DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in rCutSmart™ Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 25 units of HpyCH4III incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in rCutSmart™ Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 25 units of HpyCH4III incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in rCutSmart™ Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 25 units of HpyCH4III incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass

Assay Name/Specification	Lot # 10251735
<p>Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of Lambda DNA with HpyCH4III, ~50% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HpyCH4III.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 5 units of HpyCH4III incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) HpyCH4III is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in rCutSmart™ Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of HpyCH4III incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of HpyCH4III is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
23 Jul 2024



Michael Tonello
Packaging Quality Control Inspector
23 Jul 2024