

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Acul
Catalog Number: R0641S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10026462
Expiration Date: 10/2020
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0641S/L v2.0

Acul Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0641SVIAL	Acul	10026463	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10033230	Pass	
B7204SVIAL	CutSmart® Buffer	10031565	Pass	

Assay Name/Specification	Lot # 10026462
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 5 Units of Acul incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Acul is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 units of Acul incubated for 4 hours at 37°C results in <50% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 25 units of Acul incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	Pass



R0641S / Lot: 10026462

Page 1 of 2



Assay Name/Specification	Lot # 10026462
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with Acul, ~50% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
~75% can be recut with Acul.	

This product has been tested and shown to be in compliance with all specifications.

Stephanie Cornelio Production Scientist

25 Oct 2018

Michael Tonello

Packaging Quality Control Inspector

16 Jan 2019

