

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Pcil   |
|------------------------|--|
| Catalog Number:        | R0655S   |
| Concentration:         | 10,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Packaging Lot Number:  | 10114618   |
| Expiration Date:       | 02/2023  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 500 μg/ml BSA  |
| Specification Version: | PS-R0655S/L v1.0   |

| Pcil Component List |                              |            |                      |  |
|---------------------|------------------------------|------------|----------------------|--|
| NEB Part Number     | Component Description        | Lot Number | Individual QC Result |  |
| R0655SVIAL          | Pcil                         | 10100404   | Pass                 |  |
| B7024AVIAL          | Gel Loading Dye, Purple (6X) | 10105819   | Pass                 |  |
| B6003SVIAL          | NEBuffer™ r3.1               | 10110766   | Pass                 |  |

| Assay Name/Specification   | Lot # 10114618 |
|--|----------------|
| <b>Endonuclease Activity (Nicking)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a<br>minimum of 10 Units of Pcil incubated for 4 hours at 37°C results in <10% conversion<br>to the nicked form as determined by agarose gel electrophoresis.            | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Pcil incubated for 4<br>hours at 37°C releases <0.1% of the total radioactivity. | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 10-fold over-digestion of pXba DNA with Pcil, >95% of the DNA fragments can<br>be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95%<br>can be recut with Pcil.  | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of pXba DNA and a minimum of 50<br>Units of Pcil incubated for 16 hours at 37°C results in a DNA pattern free of  | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification  | Lot # 10114618 |
|---|----------------|
| detectable nuclease degradation as determined by agarose gel electrophoresis. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist 21 Jul 2021

Michae 1...

Michael Tonello Packaging Quality Control Inspector 21 Jul 2021

