

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Zral   |
|------------------------|--|
| Catalog Number:        | R0659S   |
| Concentration:         | 10,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Lot Number:            | 10041055   |
| Expiration Date:       | 03/2021  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 500 μg/ml BSA  |
| Specification Version: | PS-R0659S/L v1.0   |

| Zral Component List    |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0659SVIAL             | Zral                  | 10041056   | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer      | 10043912   | Pass                 |  |

| Assay Name/Specification  | Lot # 10041055 |
|---|----------------|
| Endonuclease Activity (Nicking)<br>A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled M13mp18 DNA and<br>a minimum of 10 units of Zral incubated for 4 hours at 37°C results in <20%<br>conversion to the nicked form as determined by agarose gel electrophoresis.                          | Pass           |
| Exonuclease Activity (Radioactivity Release)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Zral incubated for 4<br>hours at 37°C releases <0.1% of the total radioactivity.   | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 10-fold over-digestion of Lambda DNA with Zral, >95% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>>95% can be recut with Zral.   | Pass           |
| Non-Specific DNase Activity (16 Hour)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of<br>10 Units of Zral incubated for 16 hours at 37°C results in a DNA pattern free of<br>detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification  | Lot # 10041055 |
|---|----------------|
| Protein Purity Assay (SDS-PAGE)<br>ZraI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue<br>detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist 26 Mar 2019

 $\ge l$ 

Jay Minichiello Packaging Quality Control Inspector 13 Jun 2019

