

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: PI-PspI
Catalog Number: R0695S
Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 µg

of pAKR7 Xmnl-linearized Control Plasmid in 1 hour at 65°C in a

total reaction volume of 50 μl.

Packaging Lot Number: 10110861
Expiration Date: 06/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R0695S/L v1.0

| PI-PspI Component List | | | | |
|------------------------|--|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0695SVIAL | PI-PspI | 10110860 | Pass | |
| N0421SVIAL | pAKR7 Xmnl-linearized Control Plasmid | 10090822 | Pass | |
| B9200SVIAL | Recombinant Albumin, Molecular Biology G | 10106371 | Pass | |
| B0695SVIAL | NEBuffer™ PI-PspI | 10090821 | Pass | |

| Assay Name/Specification | Lot # 10110861 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer PI-Pspl containing 1 µg of supercoiled PhiX174 DNA and a minimum of 15 Units of PI-Pspl incubated for 4 hours at 65°C results in <20% | Pass |
| conversion to the nicked form as determined by agarose gel electrophoresis. | |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer PI-PspI containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of PI-PspI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of pAKR7-XmnI DNA with PI-PspI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with PI-PspI. | Pass |
| Non-Specific DNase Activity (16 Hour) | Pass |



R0695S / Lot: 10110861

Page 1 of 2



| Assay Name/Specification | Lot # 10110861 |
|---|----------------|
| A 50 µl reaction in NEBuffer PI-Pspl containing 1 µg of pAKR7-Xmnl DNA and a minimum | |
| of 5 Units of PI-PspI incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

23 Jun 2021

Michael Tonello

Packaging Quality Control Inspector

23 Jun 2021



R0695S / Lot: 10110861

Page 2 of 2