

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: PI-Pspl
Catalog Number: R0695S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 µg

of pAKR7 Xmnl-linearized Control Plasmid in 1 hour at 65°C in a

total reaction volume of 50 μl.

Packaging Lot Number: 10140866
Expiration Date: 11/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0695S/L v1.0

PI-PspI Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0695SVIAL	PI-PspI	10127770	Pass	
N0421SVIAL	pAKR7 Xmnl-linearized Control Plasmid	10127774	Pass	
B9200SVIAL	Recombinant Albumin, Molecular Biology G	10106371	Pass	
B0695SVIAL	NEBuffer™ PI-PspI	10127773	Pass	

Assay Name/Specification	Lot # 10140866
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer PI-PspI containing 1 μg of supercoiled PhiX174 DNA and a minimum of 15 Units of PI-PspI incubated for 4 hours at 65°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 μl reaction in NEBuffer PI-PspI containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of PI-PspI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 5-fold over-digestion of pAKR7-XmnI DNA with PI-PspI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with PI-PspI.	Pass
Non-Specific DNase Activity (16 Hour)	Pass



R0695S / Lot: 10140866

Page 1 of 2

Assay Name/Specification	Lot # 10140866
A 50 μl reaction in NEBuffer PI-Pspl containing 1 μg of pAKR7-Xmnl DNA and a minimum	
of 5 Units of PI-PspI incubated for 16 hours at 65°C results in a DNA pattern free	
of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Pengha Zhang Production Scientist

26 May 2022

Michael Tonello

Packaging Quality Control Inspector

26 May 2022



R0695S / Lot: 10140866

Page 2 of 2