

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: TspMI
Catalog Number: R0709S
Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pUCAdeno plasmid DNA in 1 hour at 75°C in a total reaction volume

of 50 μl.

Packaging Lot Number: 10058186
Expiration Date: 04/2020
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 20 mM Tris-HCl (pH 8.0), 1 mM DTT, 1 mM EDTA, 50%

Glycerol, 0.10% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0709S/L v1.0

| TspMI Component List   |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0709SVIAL             | TspMI                 | 10058187   | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer      | 10046082   | Pass                 |  |

| Assay Name/Specification   | Lot # 10058186 |
|--|----------------|
| Non-Specific DNase Activity (16 Hour)  | Pass           |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pUCAdeno DNA and a minimum of 5 Units of TspMl incubated for 16 hours at 75°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.                       |                |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pUCAdeno DNA with TspMI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >75% can be recut with TspMI.                          | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of TspMI incubated for 4 hours at 75°C releases <0.1% of the total radioactivity. | Pass           |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 Units of TspMl incubated for 4 hours at 75°C results in <10%  | Pass           |



R0709S / Lot: 10058186

Page 1 of 2



| Assay Name/Specification  | Lot # 10058186 |
|---|----------------|
| conversion to the nicked form as determined by agarose gel electrophoresis. |                |

This product has been tested and shown to be in compliance with all specifications.

**G**reen Duquett **Production Scientist** 

02 Aug 2019

Josh Hersey Packaging Quality Control Inspector

22 Oct 2019

