

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: TspMI
Catalog Number: R0709S
Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBC4 plasmid DNA in 1 hour at 75°C in a total reaction volume of

50 μl.

Packaging Lot Number: 10150948
Expiration Date: 11/2022
Storage Temperature: -20°C

Storage Conditions: 20 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 1 mM EDTA, 50% Glycerol,

0.10% Triton® X-100, 200 μg/ml BSA, (pH 8.0 @ 25C)

Specification Version: PS-R0709S/V v2.0

| TspMI Component List   |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0709SVIAL             | TspMI                 | 10150947   | Pass                 |  |
| B6004SVIAL             | rCutSmart™ Buffer     | 10150371   | Pass                 |  |

| Assay Name/Specification  | Lot # 10150948 |
|---|----------------|
| Ligation and Recutting (Terminal Integrity)  After a 10-fold over-digestion of pBC4 DNA with TspMI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, ≥75%   | Pass           |
| can be recut with TspMI.  Exonuclease Activity (Radioactivity Release)  A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and   | Pass           |
| double-stranded [ ³H] E. coli DNA and a minimum of 50 units of TspMI incubated for 4 hours at 75°C releases <0.1% of the total radioactivity.   |                |
| Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 5 units of TspMI incubated for 4 hours at 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of pBC4 DNA and a minimum of 5 units of TspMl incubated for 16 hours at 75°C results in a DNA pattern free of  | Pass           |



R0709S / Lot: 10150948

Page 1 of 2



| Assay Name/Specification  | Lot # 10150948 |
|---|----------------|
| detectable nuclease degradation as determined by agarose gel electrophoresis. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

27 May 2022

Erin Varney

Packaging Quality Control Inspector

27 May 2022



R0709S / Lot: 10150948

Page 2 of 2