

New England Biolabs Certificate of Analysis

Product Name: Psil-v2
Catalog Number: R0744S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10104505
Expiration Date: 01/2023
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)
Specification Version: PS-R0744S/L v1.0

Psil-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0744SVIAL	Psil-v2	10097868	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10091457	Pass
B6004SVIAL	rCutSmart™ Buffer	10103711	Pass

Assay Name/Specification	Lot # 10104505
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Psil-v2 is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Psil-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Psil-v2.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of Psil-v2 is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
Endonuclease Activity (Nicking)	Pass

Assay Name/Specification	Lot # 10104505
<p>A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 10 units of Psil-v2 incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	
<p>Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of Psil-v2 incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 50 units of Psil-v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Psil-v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Psil-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
13 Apr 2021



Michael Tonello
Packaging Quality Control Inspector
13 Apr 2021