

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: HindIII-HF®
Catalog Number: R3104M
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10063075
Expiration Date: 10/2021
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R3104T/M v1.0

HindIII-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3104MVIAL	HindIII-HF®	10056692	Pass	
B7204SVIAL	CutSmart® Buffer	10061304	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10046085	Pass	

Assay Name/Specification	Lot # 10063075
Protein Purity Assay (SDS-PAGE) HindIII-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 200 Units of HindIII-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in 50 μl PhiX174 NEBuffer 4 60 1 μg HindIII-HF™ PhiX174 37 60 units% conversion to the nicked form as determined by agarose gel electrophoresis. incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	Pass



R3104M / Lot: 10063075

Page 1 of 2

Assay Name/Specification	Lot # 10063075
double-stranded [³H] E. coli DNA and a minimum of 200 units of HindIII-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 100-fold over-digestion of Lambda DNA with HindIII-HF™, >95% of the DNA	
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with HindIII-HF™.	

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist

21 Oct 2019

Michael Tonello

Packaging Quality Control Inspector

22 Jan 2020