

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: HindIII-HF®
Catalog Number: R3104S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number:10014308Expiration Date:03/2020Storage Temperature:-20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R3104S/L v1.0

HindIII-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3104SVIAL	HindIII-HF®	0071803	Pass	
B7204SVIAL	CutSmart® Buffer	3091805	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	0251805	Pass	

Assay Name/Specification	Lot # 10014308
Protein Purity Assay (SDS-PAGE) HindIII-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of HindIII-HF™ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart [™] Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of HindIII-HF [™] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 200 Units of HindIII-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass



R3104S / Lot: 10014308 Page 1 of 2

Assay Name/Specification	Lot # 10014308
electrophoresis.	
Ligation and Recutting (Terminal Integrity) After a 100-fold over-digestion of Lambda DNA with HindIII-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HindIII-HF™.	Pass

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang Production Scientist

26 Jun 2018

Michael Tonello

Packaging Quality Control Inspector

26 Jun 2018



R3104S / Lot: 10014308

Page 2 of 2