

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: HindIII-HF®

Catalog Number: R3104S

Concentration: 20,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10107896
Expiration Date: 03/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R3104S/L v1.0

| HindIII-HF® Component List |                              |            |                      |  |
|----------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>     | Component Description        | Lot Number | Individual QC Result |  |
| R3104SVIAL                 | HindIII-HF®                  | 10101186   | Pass                 |  |
| B7024AVIAL                 | Gel Loading Dye, Purple (6X) | 10105817   | Pass                 |  |
| B6004SVIAL                 | rCutSmart™ Buffer            | 10108412   | Pass                 |  |

| Assay Name/Specification   | Lot # 10107896 |
|--|----------------|
| <b>Ligation and Recutting (Terminal Integrity)</b> After a 100-fold over-digestion of Lambda DNA with HindIII-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HindIII-HF™.                              | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 200 Units of HindIII-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of HindIII-HF™ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.               | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and   | Pass           |



R3104S / Lot: 10107896

Page 1 of 2

| Assay Name/Specification   | Lot # 10107896 |
|--|----------------|
| double-stranded [ ³H] E. coli DNA and a minimum of 200 units of HindIII-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. |                |
| Protein Purity Assay (SDS-PAGE) HindIII-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.                          | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

16 Jun 2021

Michael Tonello

Packaging Quality Control Inspector

16 Jun 2021



R3104S / Lot: 10107896

Page 2 of 2