

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: HindIII-HF®
Catalog Number: R3104T
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10093257
Expiration Date: 12/2022
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R3104T/M v1.0

HindIII-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3104TVIAL	HindIII-HF®	10093258	Pass	
B7204SVIAL	CutSmart® Buffer	10092681	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089404	Pass	

Assay Name/Specification	Lot # 10093257
Endonuclease Activity (Nicking) A 50 μl reaction in 50 μl PhiX174 NEBuffer 4 60 1 μg HindIII-HF™ PhiX174 37 60 units% conversion to the nicked form as determined by agarose gel electrophoresis. incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of HindIII-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 100-fold over-digestion of Lambda DNA with HindIII-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HindIII-HF™.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of Lambda DNA and a minimum of	Pass



R3104T / Lot: 10093257

Page 1 of 2

Assay Name/Specification	Lot # 10093257
200 Units of HindIII-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) HindIII-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

19 Jan 2021

Michael Tonello

Packaging Quality Control Inspector

19 Jan 2021



R3104T / Lot: 10093257

Page 2 of 2