

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Scal-HF®
Catalog Number: R3122M
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total

reaction volume of 50 µl.

Packaging Lot Number: 10159269
Expiration Date: 08/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

200 μg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R3122M v2.0

Scal-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3122MVIAL	Scal-HF®	10159268	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10156427	Pass	
B6004SVIAL	rCutSmart™ Buffer	10156428	Pass	

Assay Name/Specification	Lot # 10159269
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	
60 units of Scal-HF® incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE)	Pass
Scal-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	
detection.	
Endonuclease Activity (Nicking)	Pass
A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and	
a minimum of 20 units of Scal-HF® incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 2-fold over-digestion of Lambda DNA with Scal-HF®, >95% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	



R3122M / Lot: 10159269

Page 1 of 2

Assay Name/Specification	Lot # 10159269
>95% can be recut with Scal-HF®.	
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of Scal-HF® incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Scal-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 20 units of Scal-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

19 Aug 2022

Michael Tonello

Packaging Quality Control Inspector

19 Aug 2022



R3122M / Lot: 10159269

Page 2 of 2