

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | SspI-HF®   |
|------------------------|--|
| Catalog Number:        | R3132L   |
| Concentration:         | 20,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Packaging Lot Number:  | 10090250   |
| Expiration Date:       | 09/2022  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R3132S/L v1.0   |

| SspI-HF® Component List |                              |            |                      |  |
|-------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number         | Component Description        | Lot Number | Individual QC Result |  |
| R3132LVIAL              | SspI-HF®                     | 10082804   | Pass                 |  |
| B7204SVIAL              | CutSmart® Buffer             | 10085423   | Pass                 |  |
| B7024AVIAL              | Gel Loading Dye, Purple (6X) | 10084971   | Pass                 |  |

| Assay Name/Specification   | Lot # 10090250 |
|--|----------------|
| Exonuclease Activity (Radioactivity Release)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of SspI-HF <sup>™</sup> incubated<br>for 4 hours at 37°C releases <0.1% of the total radioactivity.    | Pass           |
| Ligation and Recutting (Terminal Integrity)<br>After a 10-fold over-digestion of Lambda DNA with SspI-HF <sup>™</sup> , >95% of the DNA<br>fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated<br>fragments, >95% can be recut with SspI-HF <sup>™</sup> .                                       | Pass           |
| Non-Specific DNase Activity (16 Hour)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of<br>100 Units of SspI-HF <sup>™</sup> incubated for 16 hours at 37°C results in a DNA pattern free<br>of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Penghua Zhang Production Scientist 05 Nov 2020

Michae

Michael Tonello Packaging Quality Control Inspector 05 Nov 2020

