

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sall-HF®
Catalog Number: R3138S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Lot Number: 10027079
Expiration Date: 08/2020
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 300 μg/ml BSA

Specification Version: PS-R3138S/L v1.0

Sall-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3138SVIAL	Sall-HF®	10020577	Pass	
B7204SVIAL	CutSmart® Buffer	10021116	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10018416	Pass	

Assay Name/Specification	Lot # 10027079
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Sall-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 Units of Sall-HF™ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of Sall-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 50-fold over-digestion of Adenovirus-2 DNA with Sall-HF™, >95% of the DNA	Pass



R3138S / Lot: 10027079

Page 1 of 2



Assay Name/Specification	Lot # 10027079
fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with Sall-HF™.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 200 Units of Sall-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist

28 Aug 2018

Michael Tonello

Packaging Quality Control Inspector

29 Oct 2018



R3138S / Lot: 10027079

Page 2 of 2