

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Sall-HF®
Catalog Number:	R3138S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10171946
Expiration Date:	08/2024
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 300 μg/ml BSA, (pH 7.5 @ 25°C)
Specification Version:	PS-R3138S/L/V v2.0

Sall-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3138SVIAL	Sall-HF®	10159234	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10165690	Pass	
B6004SVIAL	rCutSmart™ Buffer	10168650	Pass	

Assay Name/Specification	Lot # 10171946
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of Sall-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 50-fold over-digestion of pBC4XS DNA with Sall-HF®, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with Sall-HF®.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of pBR322 DNA and a minimum of 200 units of Sall-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Sall-HF®, religated	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10171946
and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Sall-HF® incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jianying Luo Production Scientist 09 Aug 2022

2. 01 Michae

Michael Tonello Packaging Quality Control Inspector 01 Dec 2022

