

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pvull-HF®
Catalog Number: R3151S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37° in a total reaction volume of 50 μl.

Packaging Lot Number: 10096685
Expiration Date: 10/2022
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R3151S/L v1.0

| Pvull-HF® Component List |                              |            |                      |  |
|--------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>   | Component Description        | Lot Number | Individual QC Result |  |
| R3151SVIAL               | Pvull-HF®                    | 10088424   | Pass                 |  |
| B7204SVIAL               | CutSmart® Buffer             | 10093117   | Pass                 |  |
| B7024AVIAL               | Gel Loading Dye, Purple (6X) | 10089405   | Pass                 |  |

| Assay Name/Specification   | Lot # 10096685 |
|--|----------------|
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of Pvull-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 Units of Pvull-HF™ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.              | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 200 units of PvuII-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.                | Pass           |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Pvull-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated   | Pass           |



R3151S / Lot: 10096685

Page 1 of 2



| Assay Name/Specification                     | Lot # 10096685 |
|--|----------------|
| fragments, >95% can be recut with PvuII-HF™. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

10 Feb 2021

Josh Hersey

Packaging Quality Control Inspector

10 Feb 2021

