

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Sphl-HF®
Catalog Number: R3182M
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10107068
Expiration Date: 05/2023
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R3182M v1.0

SphI-HF® Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3182MVIAL	SphI-HF®	10107066	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10093123	Pass	
B6004SVIAL	rCutSmart™ Buffer	10108730	Pass	

Assay Name/Specification	Lot # 10107068
Protein Purity Assay (SDS-PAGE) Sphl-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass
detection.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	Pass
100 units of SphI-HF™ incubated for 16 hours at 37°C results in a DNA pattern free	
of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)  After a 20-fold over-digestion of Lambda DNA with Sphl-HF™, >95% of the DNA	Pass
fragments can be ligated with T4 DNA ligase in 16 hours at 16 °C. Of these ligated fragments, >95% can be recut with SphI-HF™.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Sphl-HF™ incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	



R3182M / Lot: 10107068 Page 1 of 2

Assay Name/Specification	Lot # 10107068
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 60 units of Sphl-HF™ incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of SphI-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

06 Jul 2021

Josh Hersey

Packaging Quality Control Inspector

06 Jul 2021



R3182M / Lot: 10107068

Page 2 of 2