

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: NotI-HF®
Catalog Number: R3189L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10050226
Expiration Date: 06/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R3189S/L v1.0

NotI-HF® Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3189LVIAL	Notl-HF®	10046576	Pass	
B7204SVIAL	CutSmart® Buffer	10043913	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10043909	Pass	

Assay Name/Specification	Lot # 10050226
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 Units of Notl-HF™ incubated for 4 hours at 37°C results in <10%	Pass
conversion to the nicked form as determined by agarose gel electrophoresis. <b>Exonuclease Activity (Radioactivity Release)</b> A 50 μl reaction in CutSmart <sup>™</sup> Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 200 units of Notl-HF <sup>™</sup> incubated	Pass
for 4 hours at 37°C releases <0.1% of the total radioactivity.  Ligation and Recutting (Terminal Integrity)  After a 10-fold over-digestion of pBC4 DNA with Notl-HF™, >95% of the DNA fragments	Pass
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NotI-HF <sup>TM</sup> .  Non-Specific DNase Activity (16 Hour) A 50 μI reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of pBC4 DNA and a minimum of 200 Units of NotI-HF <sup>TM</sup> incubated for 16 hours at 37°C results in a DNA pattern free	Pass



R3189L / Lot: 10050226

Page 1 of 2



Assay Name/Specification	Lot # 10050226
of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Doreen Duquette Production Scientist

28 May 2019

Jay Minichiello

Packaging Quality Control Inspector

31 Jul 2019



R3189L / Lot: 10050226

Page 2 of 2