

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Notl-HF®   |
|------------------------|--|
| Catalog Number:        | R3189L   |
| Concentration:         | 20,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Packaging Lot Number:  | 10094895   |
| Expiration Date:       | 01/2023  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA   |
| Specification Version: | PS-R3189S/L v1.0   |

| NotI-HF® Component List |                              |            |                      |  |
|-------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number         | Component Description        | Lot Number | Individual QC Result |  |
| R3189LVIAL              | NotI-HF®                     | 10094892   | Pass                 |  |
| B7204SVIAL              | CutSmart® Buffer             | 10092684   | Pass                 |  |
| B7024AVIAL              | Gel Loading Dye, Purple (6X) | 10089404   | Pass                 |  |

| Assay Name/Specification  | Lot # 10094895 |
|---|----------------|
| Endonuclease Activity (Nicking)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and<br>a minimum of 100 Units of NotI-HF <sup>™</sup> incubated for 4 hours at 37°C results in <10%<br>conversion to the nicked form as determined by agarose gel electrophoresis.           | Pass           |
| Exonuclease Activity (Radioactivity Release)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of NotI-HF <sup>™</sup> incubated<br>for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass           |
| Ligation and Recutting (Terminal Integrity)<br>After a 10-fold over-digestion of pBC4 DNA with NotI-HF <sup>™</sup> , >95% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>>95% can be recut with NotI-HF <sup>™</sup> .                                      | Pass           |
| Non-Specific DNase Activity (16 Hour)<br>A 50 μl reaction in CutSmart™ Buffer containing 1 μg of pBC4 DNA and a minimum of<br>200 Units of NotI-HF™ incubated for 16 hours at 37⁰C results in a DNA pattern free  | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification   | Lot # 10094895 |
|--|----------------|
| of detectable nuclease degradation as determined by agarose gel electrophoresis. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist 26 Jan 2021

Michae 1. .

Michael Tonello Packaging Quality Control Inspector 26 Jan 2021

