

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Notl-HF® |
|------------------------|---|
| Catalog Number: | R3189M |
| Concentration: | 100,000 U/mI |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μg of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl. |
| Lot Number: | 10009920 |
| Expiration Date: | 04/2020 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA |
| Specification Version: | PS-R3189M v1.0 |

| NotI-HF® Component List | | | | |
|-------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R3189MVIAL | NotI-HF® | 0041804 | Pass | |
| B7204SVIAL | CutSmart® Buffer | 3061804 | Pass | |
| B7024SVIAL | Gel Loading Dye, Purple (6X) | 0231804 | Pass | |

| Assay Name/Specification | Lot # 10009920 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 Units of NotI-HF [™] incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of NotI-HF [™] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pBC4 DNA with NotI-HF [™] , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NotI-HF [™] . | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of pBC4 DNA and a minimum of 200 Units of NotI-HF™ incubated for 16 hours at 37⁰C results in a DNA pattern free | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10009920 |
|--|----------------|
| of detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang

Penghua Zhang Production Scientist 23 May 2018

Michae 11.

Michael Tonello Packaging Quality Control Inspector 23 May 2018

