

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	NotI-HF®
Catalog #:	R3189S/L
Concentration:	20,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
<i>Lot</i> #:	0041603
Assay Date:	03/2016
Expiration Date:	03/2018
Storage Temp:	-20°C
Storage Conditions:	50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R3189S/L v1.0
Effective Date:	14 Nov 2013

Assay Name/Specification (minimum release criteria)   Endonuclease Activity (Nicking) - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of   supercoiled PhiX174 DNA and a minimum of 100 Units of NotI-HF <sup>™</sup> incubated for 4 hours at 37°C results in   <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of pBC4 DNA with NotI-HF <sup>TM</sup> , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NotI-HF <sup>TM</sup> .	
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart <sup>TM</sup> Buffer containing 1 µg of pBC4 DNA and a minimum of 200 Units of NotI-HF <sup>TM</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

otimon

Authorized by Derek Robinson 14 Nov 2013



Joby F. Unas Inspected by

Inspected by Toby Claus 25 Mar 2016