

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Ncol-HF®
Catalog Number:	R3193M
Concentration:	100,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10091240
Expiration Date:	07/2022
Storage Temperature:	-20°C
Storage Conditions:	200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R3193M v1.0

Ncol-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3193MVIAL	Ncol-HF®	10079049	Pass	
B7204SVIAL	CutSmart® Buffer	10089399	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084973	Pass	

Assay Name/Specification	Lot # 10091240
Protein Purity Assay (SDS-PAGE) Ncol-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of Ncol-HF [™] incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with NcoI-HF [™] , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NcoI-HF [™] .	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Ncol-HF [™] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10091240
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of Ncol-HF [™] incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of Ncol-HF [™] , religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 18 Nov 2020

Much

Michael Tonello Packaging Quality Control Inspector 18 Nov 2020

