

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Eagl-HF®
Catalog Number: R3505L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10163698
Expiration Date: 02/2025
Storage Temperature: -80°C

Storage Conditions: 500 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

200 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R3505S/L v3.0

| Eagl-HF® Component List | | | | |
|-------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R3505LVIAL | Eagl-HF® | 10163699 | Pass | |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10173660 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10175290 | Pass | |

| Assay Name/Specification | Lot # 10163698 |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 100 Units of Eagl-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with EagI-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EagI-HF™. | Pass |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of Eagl-HF™ incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Blue-White Screening (Terminal Integrity) A sample of Litmus38i vector linearized with a 10-fold excess of Eagl-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment | Pass |



R3505L / Lot: 10163698 Page 1 of 2

Pass

This product has been tested and shown to be in compliance with all specifications.

EagI-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sưń Production Scientist

03 Feb 2023

detection.

Michael Tonello

Packaging Quality Control Inspector

06 Feb 2023



R3505L / Lot: 10163698

Page 2 of 2