

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Apol-HF®
Catalog Number: R3566L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37 degrees C in a total reaction volume

of 50 μL

Packaging Lot Number: 10087050 Expiration Date: 08/2022 Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

200 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R3566S/L v1.0

Apol-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3566LVIAL	Apol-HF®	10082139	Pass	
B7204SVIAL	CutSmart® Buffer	10089403	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089392	Pass	

Assay Name/Specification	Lot # 10087050	
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Apol-HF, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass	
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Apol-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass	
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of Apol-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass	
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Apol-HF, >95% of the DNA fragments	Pass	



R3566L / Lot: 10087050

Page 1 of 2

Assay Name/Specification	Lot # 10087050
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Apol-HF.	
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and a minimum of 100 units of Apol-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Apol-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

03 Dec 2020

Josh Hersey

Packaging Quality Control Inspector

03 Dec 2020



R3566L / Lot: 10087050

Page 2 of 2