

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Apol-HF®
Catalog Number:	R3566S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in 1 hour at 37 degrees C in a total reaction volume of 50 μL
Lot Number:	10050299
Expiration Date:	07/2021
Storage Temperature:	-20°C
Storage Conditions:	200 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 μg/ml BSA, (pH 7.4 @ 25°C)
Specification Version:	PS-R3566S/L v1.0

Apol-HF® Compone	nt List		
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3566SVIAL	Apol-HF®	10050301	Pass
B7204SVIAL	CutSmart® Buffer	10046087	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10043910	Pass

Assay Name/Specification	Lot # 10050299
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Apol-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and 1 μl of Apol-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Apol-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Apol-HF.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and a minimum of	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10050299
100 units of ApoI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) Apol-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Apol-HF, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis Production Scientist 23 Jul 2019

AL

Minichiello Packaging Quality Control Inspector 14 Aug 2019

