

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Mfel-HF®
Catalog Number: R3589S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10101356
Expiration Date: 03/2022
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R3589S/L v2.0

Mfel-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3589SVIAL	Mfel-HF®	10101354	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10091457	Pass	
B6004SVIAL	rCutSmart™ Buffer	10103709	Pass	

Assay Name/Specification	Lot # 10101356
Protein Purity Assay (SDS-PAGE) Mfel-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass
detection.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	Pass
60 units of Mfel-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 20 units of Mfel-HF™ incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 20 units of Mfel-HF™ incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	



R3589S / Lot: 10101356

Page 1 of 2

Assay Name/Specification	Lot # 10101356 Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Mfel-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Mfel-HF™.	
Blue-White Screening (Terminal Integrity) A sample of LITMUS38i vector linearized with a 10-fold excess of Mfel-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

gene results in <1% white colonies.

06 Apr 2021

Michael Tonello

Packaging Quality Control Inspector

06 Apr 2021



R3589S / Lot: 10101356

Page 2 of 2